

Hypoglycemic Effect of Lipoic Acid, Carnitine and *Nigella Sativa* in Diabetic Rat Model

Ragaa Hamdy Mohamed Salama

Department of Medical Biochemistry, College of Medicine,
Qassim University, KSA

Abstract

Objectives: Evaluation of therapeutic potentials of α -lipoic acid (α -LA), L-carnitine, *Nigella sativa* (*N. sativa*) or combination of them in carbohydrate and lipid metabolism of DM type I.

Methods: Rat model of diabetes was induced by single i.p injection of Streptozocin (STZ) 65 mg/kg. The rats were randomly assigned to 6 groups (G): healthy reference (HR), diabetic (DM), DM treated with α -lipoic acid, DM treated with L-carnitine, DM treated with *N. sativa*, and DM treated with combination of the 3 compounds. After 30 days from onset of diabetes, serum and tissue homogenate were obtained for evaluation of glucose metabolism as fasting blood glucose, insulin, insulin sensitivity, HOMA, C-peptide, and pyruvate dehydrogenase (PDH) activity. For lipid metabolism evaluation, total cholesterol and triacylglycerol (TG) were determined. Markers of antioxidants and oxidative status as total antioxidant capacity (TAC), glutathione-S-transferase (GST), 8-hydroxy-2-deoxyguanosine (8-OH-dG) were measured.

Results: Either α -LA or *N. sativa* significantly reduced the elevated blood glucose level. The combination of 3 compounds significantly increased the level of insulin and C-peptide. Also, increased the antioxidant activity measured by TAC and decreased the oxidative damage of DNA as measured by 8-OH-dG. HOMA- β increased in G3 and G6 compared to G2. However, the decrease in TG, and total cholesterol levels were non-significant in all groups.

Conclusion: Combination of α -LA, L-carnitine and *N. sativa* will contribute significantly in improvement of the carbohydrate metabolism and to less extent lipid metabolism in diabetic rats, thus increasing the rate of success in management of DM. Also, this combination will have implications in clinical studies and clinical applications.

Key words: lipoic acid - carnitine- *Nigella sativa*- DM

Correspondence:

Ragaa H.M. Salama

Associate Professor of Medical Biochemistry
College of Medicine, Qassim University, Qassim
Tel: +966-3800050 Ext- 6043,
Fax: +966-3802082

Introduction

The overall prevalence of DM in adults in KSA is 23.7% in the age group of 30-70-years between 1995 and 2000. ⁽¹⁾ According to reports by the American Diabetes Association, people living in Saudi Arabia and the Gulf could face a lifetime risk as high as 60 per cent of developing diabetes. ⁽²⁾

α -LA is unique natural antioxidants and plays a fundamental role in the metabolism. ⁽³⁾ It is relatively safe, even in patients with renal and liver failure. It targets cellular signal transduction pathways which increased glucose uptake and utilization, antagonizing the oxidative and inflammatory stresses. ⁽⁴⁾ Acetyl-L-carnitine had a prophylactic and therapeutic potential on the progressive diabetic neuropathy. ⁽⁵⁾ L-carnitine acts as a carrier for long-chain fatty acids from the cytosol into mitochondria for β -oxidation, hence sustains energy supply. Combination of α -LA, acetyl-L-carnitine, nicotinamide, and biotin effectively improved glucose tolerance, decreased the level of circulating free fatty acid, and prevented the reduction of mitochondrial biogenesis in skeletal muscle of diabetic rats. ⁽⁶⁾ Acetyl-L-carnitine safely ameliorated arterial hypertension, insulin resistance, impaired glucose tolerance, and hypoadiponectinemia in DM. ⁽⁷⁾ Several studies showed that extracts from the seeds of *N. sativa* have antidiabetic effects and may be used as a treatment for diabetes due to their stimulatory effect on beta cell function with consequent increase in serum insulin level. ⁽⁸⁾

The aim of this study is to evaluate the isolated effect of α -LA, L-carnitine and *N. sativa* oil on carbohydrate and lipid metabolism of diabetic rats. Also, to investigate the therapeutic potential of combination of the three compounds, hence, improving the rate of success in management of DM.

Materials and Methods

Chemicals and Experimental Design

Streptozocin (STZ) (EC242-646-8), dihydro lipoic acid (α -LA), (T8260), and L-carnitine hydrochloride (EC229) purchased from sigma. *N. sativa* oil purchased from Pharco-Company, Egypt.

The study carried out on healthy adult male Sprague-Dawley rats weighing 150-200g. Their

ages ranged from 15-20 weeks. The rats housed conventionally in clean cages under a 12-hour light-dark cycle at 37°C and fed with standard food and water *ad libitum*. The experimental and feeding protocols of the animals approved and performed according to the guidelines of Animal House and ethical standards of Faculty of Medicine, Qassim University, KSA. The experiment carried out on 6 groups. The first group (7 rats) was the healthy reference (H.R) group, injected once with 1 ml saline i.p. The second group (G2) was the diabetic rat model. Rats injected once with 65 mg/kg STZ i.p following overnight fasting. ⁽⁹⁾ After 3 days, 42 rats that developed diabetes, (sever polyurea, and fasting blood glucose over 200 mg/dl) divided randomly into 4 subgroups received daily oral treatment in the form of α -LA (100 mg/kg) for group (3), ⁽¹⁰⁾ L-carnitine (100 mg/kg) for group (4), *N. sativa* oil (500 mg/kg) for group (5) or combination of the 3 compounds for group (6) in the same dose like the above groups for 30 days. The animal fasted overnight, then weighted, blood samples obtained, and sacrificed. Pancreas, kidney, and liver removed, then homogenized separately, in 2,5,20 ml ice cold PBS, respectively, centrifuged at 4000xg for 10 min at 4°C and supernatant kept at -70°C for further biochemical measurements.

Biochemical Assays:

Insulin and C-peptide ELISA kits purchased from DRG, Germany. Glucose determined using the glucose oxidase – peroxidase – chromogen from bioMérieux® SA, France. The total antioxidant capacity (TAC), 8-hydroxy-2-deoxyGuanosine (8-OH-dG) and Glutathione S-transferase (GST) kits purchased from Cayman's Chemical Company, USA. TAC measured the combined antioxidant activities of all constituents. 8-OH-dG kit quantified both free 8-OH-dG and DNA-incorporated 8-OH-dG. GST assay kit measured the total GST activity (cytosolic and microsomal). Pyruvate dehydrogenase (PDH) enzyme activity determined by kit (MSP18) from MitoSciences. Total protein measured by available commercial kit. Two types of HOMA calculated, HOMA-insulin resistance (HOMA-IR) and HOMA - β -cell function (HOMA- β) which is % β of β -cell function. ⁽¹¹⁾ Triglyceride

glucose index (TyG index) which is a simple measure of insulin sensitivity.⁽¹²⁾

SPSS version 16 for windows used for statistical analysis. Data are expressed as means \pm SEM, (n = 7). One way analysis of variance (ANOVA) followed by LSD to judge the difference between different groups. ROC curves, Pearson correlation coefficient (r) used to detect the relation between markers. Significance level was accepted at $p \leq 0.05$.

Results

In this study, the diabetic rat model (G2) developed hyperglycemia that negatively correlate with insulin level, C-peptide and TAC ($r = -0.5, -0.6, -0.6$; $p=0.01, 0.001, 0.001$, respectively) that clearly explain one of the possible mechanisms of DM type I. There is significant decrease level of glucose in G3, G5, G6 ($p<0.05$) compared to G2; however, in G4 (DM+ carnitine), there is significant increase in glucose level of hepatic homogenate (Figure 1) that may be due to increase rate of β -oxidation and gluconeogenesis.




Figure 1: Glucose level in tissue homogenates




Figure 2: Insulin level in tissue homogenates

In figure 2, G2 showed decreased insulin level of pancreatic and renal homogenate while in liver homogenate increased, but, it reached near normal level in pancreatic tissue of treated groups (G3-G6).




Figure 3: C-peptide level in tissue homogenates




Figure 4: Pyruvate dehydrogenase level in tissue homogenates

There is increase in c-peptide level in pancreatic tissue homogenate in treated groups (G3-G6) but, the highest increment in G6 compared to G2 (Figure 3). In Figure 4, significant increased level of PDH activity in liver of DM (G2) and (G5) (DM+*N. sativa*), kidney homogenate of G3 (DM + α -LA). That is because; α -LA is essential cofactor for mitochondrial enzymes, including pyruvate dehydrogenase and α -ketoglutarate dehydrogenase. Figure 5 showed decreased level of TAC in pancreatic and renal homogenates of G2 compared to G1, while the other groups (G3-G6) showed increased level of TAC

except pancreatic homogenate of G6 was decreased. Figure 6 showed normalization of GST level in various organs of different groups (except G5). There is significant increase in 8-OH-dG in pancreas of G2 compared to G1, and significant decrease in pancreatic homogenate of G3 and renal homogenate of G4 (Figure 7). A positive correlation between 8-OH-dG and blood glucose level ($r = 0.45$, $p=0.05$) and negative correlation with s. insulin level ($r = -0.6$, $p=0.01$) were detected. That can explain hyperglycemia due to decrease insulin level leads to increase DNA oxidative damage.




Figure 5: Total antioxidant level in tissue homogenates




Figure 6: Glutathione –S- transferase level in tissue homogenates




Figure 7: DNA oxidative damage level in tissue homogenates

Table1: Biochemical markers of glucose homeostasis in serum of diabetic mice and control groups

| | G1 | G2 | G3 | G4 | G5 | G6 |
|-------------------------------|--------------|----------------|-------------|--------------|---------------|--------------|
| Glucose mg/dl | 78.17 ± 7.97 | 247.31 ± 11.74 | 181. 1±1 .2 | 182.1 ± 7.58 | 191.17 ± 1.13 | 164.61± 1. 5 |
| p | 0.000 | | 0.000 | 0.000 | 0.001 | 0.000 |
| Insulin mIU/ml | 4.2 ± 1.8 | 0.5±0.01 | 3.01±0.1 | 2.1 ± 0.3 | 1.17± 0.1 | 3.75±0.25 |
| p | 0.01 | | NS | NS | NS | 0.04 |
| C-peptide ng/ml | 0.42±0.15 | 0.08±0.01 | 0.4±0.01 | 0.2±0.05 | 0.2±0.05 | 0.35± 0. 05 |
| p | 0.001 | | 0.006 | 0.04 | NS | 0.01 |
| Pyruvate dehydrogenase mg/min | 0.04±0.004 | 0.04±0.002 | 0.05±0.002 | 0.09±0.04 | 0.03±0.002 | 0.04±0.007 |
| p | NS | | NS | NS | NS | NS |
| HOMA-IR | 0.80 | 0.3 | 1.35 | 0.94 | 0.55 | 1.52 |
| HOMA- β (insulin) | 99.67% | 0.97% | 9.18% | 6.35% | 3.29% | 13.29% |
| HOMA- β (C-peptide) | 9.96% | 0.15% | 1.22% | 0.60% | 0.56% | 1.25% |

Group (1) healthy reference (HR), group (2) received (STZ) only; diabetic rat model. Group (3) received lipoic acid (100 mg/kg orally) in diabetic rat, daily, group (4) received L-carnitine daily (100 mg/kg orally) in diabetic rat, group (5) received *N. sativa* oil daily (500 mg/kg orally) in diabetic rat, group (6) received daily combination of, lipoic acid, carnitine and *N. sativa* oil in diabetic rat. ANOVA test followed by LSD for the significant values ($p > 0.05$), compared all groups to group 2. HOMA-IR i.e Homeostatic Model Assessment-insulin resistance, HOMA- β i.e Homeostatic Model Assessment %β of β-cell function.

Table 2: Oxidative and antioxidant status in serum of diabetic mice and control groups

| | G1 | G2 | G3 | G4 | G5 | G6 |
|--------------------------|------------|-------------|-----------|-------------|-------------|-------------|
| TAC mM | 1.27± 0.10 | 0.66 ±0 .07 | 1.1± 0.01 | 0.79 ± 0.09 | 0.73± 0.19 | 1.07±0.01 |
| p | 0.002 | | 0.03 | NS | NS | 0.04 |
| GST activity nmol/min/ml | 3.67± 0.29 | 9.57± 0.35 | 4.79±0.57 | 7.13± 1.01 | 6.57 ± 0.87 | 4.83 ± 0.18 |
| p | 0.002 | | 0.01 | NS | NS | 0.01 |
| 8-OH-G pg/ml | 33.67±8.57 | 57.12±7.68 | 31.5±3.5 | 37.14±2.8 | 35.67±8.78 | 30.09±2.1 |
| p | 0.01 | | 0.02 | 0.01 | 0.02 | 0.01 |

Group (1) healthy reference (HR), group (2) received (STZ) only; diabetic rat model. Group (3) received lipoic acid (100 mg/kg orally) daily in diabetic rat, group (4) received L-carnitine daily (100 mg/kg orally) in diabetic rat, group (5) received *N. sativa* oil daily (500 mg/kg orally) in diabetic rat, group (6) received daily combination of the 3 compounds. TAC; total antioxidant capacity, GST; glutathione –S-tranferase 8-OH-dG; 8-hydroxy-2-deoxyguanosine.

Table 3: Biochemical markers of lipid metabolism in serum of diabetic mice and control groups

| | G1 | G2 | G3 | G4 | G5 | G6 |
|---------------------------|------------|-------------|---------------|---------------|-------------|------------|
| Total cholesterol (mg/dl) | 86.27±22.4 | 98.33± 10.7 | 80.35±0.1 | 101.04 ± 7.4 | 88.43± 6. 8 | 105.5± 6.5 |
| P | NS | | NS | NS | NS | NS |
| Triacylglycerol (mg/dl) | 90.3± 10.9 | 117.5 ± 5.9 | 100.75 ± 24.2 | 108.19 ± 12.6 | 137.4± 32.1 | 125.5± 1.5 |
| P | NS | | NS | NS | NS | NS |
| TyG index | 3529.37 | 14529.46 | 9122.91 | 9850.69 | 13133.37 | 10329.27 |

Group (1) healthy reference (HR), group (2) received (STZ) only, Group (3) received lipoic acid (100 mg/kg orally) daily in diabetic rat, group (4) received L-carnitine daily (100 mg/kg orally) in diabetic rat, group (5) received *N. sativa* oil daily (500 mg/kg orally) in diabetic rat, group (6) received daily combination of the 3 compounds. TyG i.e triglycerides and glucose index.

Discussion

In this study, the diabetic rat model (G2) showed significant increase in glucose level while insulin, c-peptide and PDH decreased (Table 1, Fig.3-6), also, TAC decreased (Table 2, Fig.5), while, GST and 8-OH-dG increased (Table 2, Fig.6,7), this could be explained by the body response to overcome the production of reactive oxygen species and/ or increased insulin degradation by glutathione insulin transhydrogenases enzyme which used reduced glutathione and GST enzyme. ⁽¹³⁾ 8-OH-dG increased especially in pancreatic homogenates (Fig. 7) indicating destruction of pancreatic tissue. HOMA-IR is decreased because this is type I DM not Type II where insulin resistance decrease but, % β of β -cell function is very low due destruction of β cell Table (1) and the triglycerides and glucose index was the highest in all groups indicate impaired lipid and carbohydrate metabolism Table (3).

On the other hand, G3 that received α -LA showed significant decrease in blood glucose, increase in insulin, c-peptide and PDH activity (Table1). In organs homogenate, there is significant decrease ($p=0.05$) in blood glucose (Fig.1) while there is non significant increase in insulin level, c-peptide, PDH activity and TAC (Fig.2-5) but with changes in GST level (Fig.6). However, there is significant decrease in 8-OH-dG in pancreatic tissue homogenate (Fig. 7). The changes in TG, and total cholesterol levels were non-significant in all groups (Table 3). Also, HOMA- β is increased in G3 compared to G2. In another studies α -LA treatment induced a significant decrease in glucose, TG, and total cholesterol levels and significant increase in serum insulin and leptin levels in diabetic animal. ⁽¹⁴⁾ The hypoglycemic and beta cell protective effects of LA occurred most probably by stimulating glucose transport protein subtype 4 (GLUT4) membrane translocation. LA is referred to as an insulin mimetic agent; hence, it is a potential new anti-diabetic agent. ⁽¹⁵⁾ For the treatment of diabetes, the recommended dosage of LA is 300–600 mg daily but for general antioxidant support, the dosage is 20–50 mg daily. Intravenous and oral LA approved for the

treatment of diabetic neuropathy in Germany. LA speeds the removal of glucose from the bloodstream at least partly by enhancing insulin function, and reduces insulin resistance. ⁽¹⁶⁾ When LA, 400mg/day was given to NIDDM patients over a period of 4 weeks with oral hypoglycemic drugs, it reduced significantly the concentration of ROS and improve HDL-C value, especially in men with metabolic syndrome. ⁽¹⁷⁾ In disagreement with this study that LA-supplementation corrected the dyslipidemia by increasing HMG-CoA reductase activity and increase lipoprotein lipase and lecithin cholesterol acyl transferase (LCAT). ⁽¹⁸⁾

During evaluation of lipid metabolism in children with DM, they found decreased plasma free carnitine level and increased fatty acids. That had negative effects on pancreatic beta-cell functions, and enhanced mitochondrial transportation of fatty acids. ⁽¹⁹⁾ L-carnitine promotes the improvement of some aspects of the diabetes such as body weight loss and glycemic level after fasting. ⁽²⁰⁾ That is why we tried carnitine in DM, in this study. G4 showed significantly decrease of glucose level, increase in insulin, c-peptide, PDH (Table 1), but it decreased significantly the level of 8-OH-dG (Table 2). There is no effect of L-carnitine on cholesterol and TG levels as shown in table 3, Here in, administration of L-carnitine alone lead to increase in glucose level in liver homogenate (Fig. 1). This could be explained by excessive hepatic glucose production through the gluconeogenesis in diabetic patients. ⁽²¹⁾ Also, increased insulin level in pancreas and kidney homogenate (Fig. 2), with increase c-peptide in pancreas (Fig.3). TAC increased in pancreatic and renal homogenates (Fig.5), but, there are non-significant changes in GST (Fig.6). Also, there is increase of 8-OH-dG in pancreatic tissue, while decrease in renal tissue (Fig.7). In agreement with our study, L-Carnitine orally administered for a period of 4 weeks did not modify insulin sensitivity or the lipid profile. ⁽²²⁾ There is a remarkable abnormality in lipid and carnitine metabolism in Korean diabetic patients. Both exercise training and supplementation of carnitine and antioxidants

improved lipid profiles and carnitine metabolism in human. ⁽²³⁾ Also, Creatine, choline and carnitine decreased in diabetic mice. ⁽²⁴⁾ Therapies for diabetic neuropathy may include replacement of acetyl-L-carnitine and replenishment of C-peptide in type 1 diabetic neuropathy. ⁽²⁵⁾

Thymoquinone at 80mg/kg b.w is associated with beneficial changes in hepatic enzyme activities, the altered activities of carbohydrate metabolic enzymes were restored to near normal, and thereby exerts potential antihyperglycemic effects. ⁽²⁶⁾ Treatment of diabetic rats with *N. sativa* extract and oil significantly decreased the serum glucose and significantly increased serum insulin ⁽²⁷⁾ hence, the designed fifth group received *N. sativa*. There was significant decrease in glucose level while insulin and c-peptide increased (Table 1). In organs homogenate, glucose level was significantly decreased (Fig.1), while insulin, c-peptide PDH and TAC increased (Fig.2-5). There was decrease of GST with variable changes in 8-OH-dG (Fig.6).

G6 that received combination of the 3 compounds showed significant decrease in glucose level while, insulin level, c-peptide, PDH and TAC were significantly increased (Table 1, 3). While, GST and 8-OH-dG decrease as shown in table 3 due to decrease degradation of insulin hormone by insulinase enzyme that use GST. These finding suggested that the combination of LA, carnitine and *N. sativa* oil have a potential therapeutic role in treatment of DM. There is significant negative correlation between glucose level, insulin and C-peptide ($r = -0.5, -0.6; p=0.05, 0.01$, respectively). % β cell function is the highest one in all treated groups (Table1). Moreover, LA, carnitine, or *N. sativa* has been used separately in treatment DM and/or its complications ^(6-8, 16, 17, 23, 25-27)

Acknowledgment:

The author acknowledges the Scientific Research Deanship, Qassim University, KSA, for the financial support of this work.

References:

1. Al-Nozha, M. M., Al-Maatouq, M. A., Al-Mazrou, Y. Y., Al-Harathi, S. S., Arafah, M. R., Khalil, M. Z., Khan, N. B., Al-Khadra, A., Al-Marzouki, K., Nouh, M. S., Moheeb Abdullah, Omer Attas, Al-Shahid, M. S., Al-Mobeireek, A. Diabetes mellitus in Saudi Arabia. Saudi Med J. 2004;25(11):1603-10.
2. <http://www.diabetes.co.uk/news/2008/Mar/60-per-cent-risk-of-diabetes-in-Saudi-Arabia.html>.
3. Singh and Jialal Singh U, Jialal I.: Alpha-lipoic acid supplementation and diabetes. (2008) Nutr Rev Nov;66(11):646-57
4. Poh Z, Goh KP. A Current Update on the Use of Alpha Lipoic Acid in the Management of Type 2 Diabetes Mellitus. Endocr Metab Immune Disord Drug Targets. 2009 Dec 1. [Epub ahead of print]
5. Ohsawa M, Miyata S, Carlsson A, Kamei J (2008). Preventive effect of acetyl-L-carnitine on the thermal hypoalgesia in streptozotocin-induced diabetic mice. Eur J Pharmacol. 7;588(2-3):213-6.
6. Shen W, Hao J, Tian C, Ren J, Yang L, Li X, Luo C, Cotman CW, Liu J. (2008): A combination of nutriment improves mitochondrial biogenesis and function in skeletal muscle of type 2 diabetic Goto-Kakizaki rats. PLoS ONE. 4; 3(6):e2328.
7. Ruggenenti P, Cattaneo D, Loriga G, Ledda F, Motterlini N, Gherardi G, Orisio S, Remuzzi G. Ameliorating hypertension and insulin resistance in subjects at increased cardiovascular risk: effects of acetyl-L-carnitine therapy. Hypertension. 2009;54(3):462-4.
8. Benhaddou-Andaloussi A, Martineau LC, Vallerand D, Haddad Y, Afshar A, Settaf A, Haddad PS. Multiple molecular targets underlie the antidiabetic effect of *Nigella sativa* seed extract in skeletal muscle, adipocyte and liver cells. Diabetes Obes Metab. 2010;12(2):148-57.
9. Kim YD, Eom SY, Zhang YW, Kim H, Park JD, Yu SD, Lee CH, Arashidani K, Kawamoto T, Kim H. Modification of the relationship between urinary 8-OHdG and hippuric acid concentration by GSTM1, GSTT1, and ALDH2 genotypes. Hum Exp Toxicol. 2010 May 20. [Epub ahead of print]

10. Chun-jun L, Qiu-mei Z, Ming-zhen L, Jing-yun Z, Pei Y, De-min Y. Attenuation of myocardial apoptosis by alpha-lipoic acid through suppression of mitochondrial oxidative stress to reduce diabetic cardiomyopathy. *Chin Med J* 2009;122(21):2580-2586
11. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and B-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985; 28: 412-9.
12. Guerrero-Romero, F, Simental-Mendía, L.E., González-Ortiz, M.c, Martínez-Abundis, E., Ramos-Zavala, M.G., Hernández-González, S.O., Jacques-Camarena, O., Rodríguez-Morán, M. The product of triglycerides and glucose, a simple measure of insulin sensitivity. Comparison with the euglycemic-hyperinsulinemic clamp. *Journal of Clinical Endocrinology and Metabolism* 2010; 95(7): 3347-3351.
13. Ansoorgep S, Rohley P, Kirschke H, Langnerb I, Wiederanders B, Hanson H. Metabolism of insulin and glucagon; glutathione-insulin transhydrogenase from microsomes of rat liver. *Eur. J. Biochem*. 1973; 32, 27-35.
14. Kandeil MA, Amin KA, Hassanin KA, Ali KM, Mohammed ET. Role of lipoic acid on insulin resistance and leptin in experimentally diabetic rats. *J Diabetes Complications*. 2011; 25(1):31-8.
15. Zhang Z, Jiang J, Yu P, Zeng X, Larrick JW, Wang Y. Hypoglycemic and beta cell protective effects of andrographolide analogue for diabetes treatment. *J Transl Med*. 2009; 16; 7:62.
16. Singh U, Jialal I: Alpha-lipoic acid supplementation and diabetes. *Nutr Rev* 2008; 66(11):646-57.
17. Gianturco V, Bellomo A, D'Ottavio E, Formosa V, Iori A, Mancinella M, Troisi G, Marigliano V. Impact of therapy with alpha-lipoic acid (ALA) on the oxidative stress in the controlled NIDDM: a possible preventive way against the organ dysfunction? *Arch Gerontol Geriatr*. 2009;49 Suppl 1:129-33.
18. Thirunavukkarasu V, Anitha Nandhini AT, Anuradha CV, Effect of alpha-lipoic acids on lipid profile in rats fed a high-fructose diet, *Exp Diabetes Res*. 2004; 5(3):195–200.
19. Coker M, Coker C, Darcan S, Can S, Orbak Z, Gökşen D. Carnitine metabolism in diabetes mellitus. *J Pediatr Endocrinol Metab*. 2002; 15(6):841-9.
20. Defani MA, Zanoni JN, Natali MR, Bazotte RB, de Miranda-Neto MH. Effect of acetyl-L-carnitine on VIP-ergic neurons in the jejunum submucous plexus of diabetic rats. *Arq Neuropsiquiatr*. 2003; 61(4):962-7. Epub 2004 Jan 6.
21. Nagashima T, Shigematsu N, Maruki R, Urano Y, Tanaka H, Shimaya A, Shimokawa T, Shibasaki M. Discovery of novel Foxo1 inhibitors for treating type 2 diabetes: Improvement of fasting glycemia in diabetic db/db mice. *Mol Pharmacol*. 2010; 24. [Epub ahead of print].
22. González-Ortiz M, Hernández-González SO, Hernández-Salazar E, Martínez-Abundis E. Effect of oral L-carnitine administration on insulin sensitivity and lipid profile in type 2 diabetes mellitus patients. *Ann Nutr Metab*. 2008;52(4):335-8
23. Cha YS. Effects of L-carnitine on obesity, diabetes, and as an ergogenic aid. *Asia Pac J Clin Nutr*. 2008; 17 Suppl 1:306-8.
24. Mallidis C, Green BD, Rogers D, Agbaje IM, Hollis J, Migaud M, Amigues E, McClure N, Browne RA. Metabolic profile changes in the testes of mice with streptozotocin-induced type 1 diabetes mellitus. *Int J Androl*. 2009 ;32(2):156-65
25. Sima AA. The heterogeneity of diabetic neuropathy. *Front Biosci*. 2008 1;(130: 4809-16
26. Pari L, Sankaranarayanan C. Beneficial effects of thymoquinone on hepatic key enzymes in streptozotocin-nicotinamide induced diabetic rats. *Life Sci*. 2009; 16; 85(23-26):830-4.
27. Abdelmeguid NE, Fakhoury R, Kamal SM, Al Wafai RJ. Effects of Nigella sativa and thymoquinone on biochemical and subcellular changes in pancreatic β -cells of streptozotocin-induced diabetic rats. *J Diabetes*. 2010;2(4):256-66

Ragaa Hamdy Mohamed Salama